

ADVANCED SOLAR RESOURCE RISK MANAGEMENT:

Solar radiation long term scenarios

Sandro Calmanti, Alessandro Dell'Aquila

ENEA



CLIM-RUN



Keywords: Solar radiation, Renewable Energy, Investments

Target groups

- Energy
- Professional organisations and federations
- Regional authorities

Relevance to the case-study requirements

Solar radiation at the Earth surface is a key parameter for climate monitoring and drives many natural processes, whose understanding and modeling can benefit from an accurate characterization of the variability of solar resource (e.g. agriculture, forestry, meteorology oceanography). In addition, the assessment of incoming solar energy is essential for applications aimed at converting solar energy into electricity. Since the renewable energy market is expanding, the prediction of the availability of resources is becoming an increasingly urgent issue for the quantification of economic losses and limits the diffusion of the exploitation of renewable energy.

The approach

Regional Climate Models (RCMs) produce high-resolution (about 20 km) climate scenarios over selected areas by taking the input at the lateral boundaries from coarser resolution (about 100 km) Global Climate Models (GCMs). RCMs enhance the quality of climate projections with respect to GCMs, especially in the presence of complex orography (Artale et al., 2010) and in the proximity of coastal areas (Feser et al., 2011). Under CLIMRUN, we have evaluated output modelling of today’s largest and most consolidated ensemble of RCM simulations produced during the EU-FP6 project ENSEMBLES over Euro-Mediterranean region (Christensen et al., 2009).

The table shows the GCMs-RCMs combinations that have been extracted from the ENSEMBLES archive to develop the CLIMRUN products on solar radiation scenarios.

		Global Model					
		HadCM3Q16	ARPEGE	BCM	ECHAM5-MPIOM r3	MIROC3.2 hires	HadCM3Q0
Regional Model	C4IRCA3						
	CNRM-RM4.5						
	DMI-HIRAM5						
	ETHZ-CLM						
	ICTP-RegCM3						
	KNMI-RACMO2						
	METNO-HIRAM						
	METO-HC HadRM3Q0						
	MPI-M-REMO						
	SMHIRCA						
	UCLM-PROMES						

References:

Artale et al., 2010. An atmosphere-ocean regional climate model for the Mediterranean area: assessment of a present climate simulation Clim. Dyn. doi:10.1007/s00382-009-0691-8


Feser et al., 2011. Regional climate models add value to global model data”. Bull. Of the American Meteorological Society 92. 1181-1192.

van der Linden P., and J.F.B. Mitchell , 2009. ENSEMBLES: Climate Change and its Impacts: Summary of research and results from the ENSEMBLES project. Met Office Hadley Centre, FitzRoy Road, Exeter EX1 3PB, UK. 160pp

# ADVANCED SOLAR RESOURCE RISK MANAGEMENT: Solar radiation long term scenarios

## The product


A1B RCMs solar radiation changes 2021-2050 vs 1961-1990; DJF



Mean change in surface downward solar radiation (W/m2) projected by A1B greenhouse gas emissions scenario from ENSEMBLES for winter (top) and summer (below).


Colours represent the average long-term change in wind speed projected for 2021-2050 with respect to 1961-1990.

A1B RCMs solar radiation changes 2021-2050 vs 1961-1990; JJA



Model data are produced at an horizontal resolution of about 25 km.

Stippled areas represent areas where more than 66% of the 16 models agree in the sign of the long-term change. Red indicates an increase, blue a decrease. We report only changes statistically significant at the 90% confidence level (t-test). The solid lines are for zero contour.



## Making the product usable

The CLIM-RUN team is seeking collaborations with stakeholders in the energy sector to make climate products usable. Areas for potential cooperation include:

- Derive tailored indices based on wind speed products
- Statistics of extremes
- Critical thresholds
- Calibrate climate models to specific sites.

*This information sheet was developed in the framework of the CLIM-RUN FP7 EU project. The product it describes should not be used without acknowledging the project and, particularly for any operational use, interaction with the authors is welcome and strongly encouraged.*